

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

COOPERS®
PANACUR® 25

ORAL ANTHELMINTIC FOR SHEEP, CATTLE AND GOATS



**ACTIVE CONSTITUENT: 25g/L
FENBENDAZOLE**

Ready to use broad-spectrum worm drench for control of benzimidazole sensitive mature and immature roundworms and lungworm in sheep, cattle and goats and to aid in control of tapeworm in sheep, cattle and goats.



F-LA3001144/03

COOPERS

1 Litre

COOPERS[®] PANACUR[®] 25

ORAL ANTHELMINTIC FOR SHEEP, CATTLE AND GOATS

READ ENCLOSED DIRECTIONS BEFORE USE

Sheep and Goats: COOPERS PANACUR 25 is indicated for the control of the following fenbendazole sensitive worms including those resistant to levamisole/morantel: **Barbers Pole Worm** (*Haemonchus contortus*); **Black Scour Worm** (*Trichostrongylus* spp.); **Large Mouthed Bowel Worm** (*Chabertia ovina*); **Nodule Worm** (*Oesophagostomum* spp.); **Thin Necked Intestinal Worm** (*Nematodirus* spp.); **Small Brown Stomach Worm** (*Teladorsagia* spp.); **Small Intestinal Worm** (*Cooperia* spp.); **Large Lung Worm** (*Dictyocaulus filaria*). COOPERS PANACUR 25 also aids in the control of **Whipworm** (*Trichuris ovis*); **Tapeworm** (*Moniezia expansa*).

Cattle: COOPERS PANACUR 25 is indicated for the control of **Barbers Pole Worm** (*Haemonchus* spp.); **Small Brown Stomach Worm** (*Ostertagia* spp.); **Stomach Hair Worm** (*Trichostrongylus axei*); **Small Intestinal Worm** (*Cooperia* spp.); **Thin Necked Intestinal Worm** (*Nematodirus* spp.); **Nodule Worm** (*Oesophagostomum radiatum*); **Whipworm** (*Trichuris ovis*); **Hookworm** (*Bunostomum phlebotomum*); **Capillaria** spp.; **Large Lungworm** (*Dictyocaulus viviparus*).

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Ph 13 11 26.

Additional Information is listed in the Material Safety Data Sheet.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

Dispose of empty container by wrapping with paper and putting in garbage.

COOPERS ANIMAL HEALTH

A division of Intervet Australia Pty Ltd
91-105 Harpin Street, Bendigo East, Vic 3550

STORE below 30°C (Room Temperature). Do not freeze. Store in tightly closed original container in a well-ventilated area. Do not store for prolonged periods in direct sunlight.

APVMA Approval No. 37097/1L/0109

FOR BATCH AND EXPIRY SEE BELOW

- ® Registered Trademarks
- ® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.
- © 2009

B-LA3001153/04



TO OPEN PEEL HERE ↘

DIRECTIONS FOR USE (ALL States)

Shake container before use: COOPERS PANACUR 25 is ready to use, no dilution or mixing is necessary. COOPERS PANACUR 25 may be administered orally through standard drenching equipment. Wash equipment after use in clean water.

Restraint

DO NOT USE on lactating or pregnant ewes where milk or milk products may be used for human consumption.

RECOMMENDATIONS FOR USE

Sheep and Goats: COOPERS PANACUR 25 is indicated for the control of the following fenbendazole sensitive worms including those resistant to levamisole/morantel: **Barbers Pole Worm** (*Haemonchus contortus*); **Black Scour Worm** (*Trichostrongylus* spp.); **Large Mouthed Bowel Worm** (*Chabertia ovina*); **Nodule Worm** (*Oesophagostomum* spp.); **Thin Necked Intestinal Worm** (*Nematodirus* spp.); **Small Brown Stomach Worm** (*Teladorsagia* spp.); **Small Intestinal Worm** (*Cooperia* spp.); **Large Lung Worm** (*Dictyocaulus filaria*). COOPERS PANACUR 25 also aids in the control of **Whipworm** (*Trichuris ovis*); **Tapeworm** (*Moniezia expansa*).

DOSE RATE: SHEEP & GOATS

Liveweight (kg)	Dose Volume (mL)	Doses per 1L Pack
10	2	500
20	4	250
30	6	167
40	8	125
50	10	100
60	12	83
70	14	72
75	15	67

1mL COOPERS PANACUR 25 per 5kg bodyweight (5mg fenbendazole/kg bodyweight). Over 75kg dose at rate of 1mL per 5kg bodyweight. Dose the mob to the heaviest animal by liveweight in each group (ewes, rams, lambs). Do not underdose. A representative sample of animals should be weighed before treatment.

Cattle: COOPERS PANACUR 25 is indicated for the control of **Barbers Pole Worm** (*Haemonchus* spp.); **Small Brown Stomach Worm** (*Ostertagia* spp.); **Stomach Hair Worm** (*Trichostrongylus axei*); **Small Intestinal Worm** (*Cooperia* spp.); **Thin Necked Intestinal Worm** (*Nematodirus* spp.); **Nodule Worm** (*Oesophagostomum radiatum*); **Whipworm** (*Trichuris ovis*); **Hookworm** (*Bunostomum phlebotomum*); **Capillaria** spp; **Large Lungworm** (*Dictyocaulus viviparus*).

DOSE RATE: CATTLE

15mL COOPERS PANACUR 25 per 50kg bodyweight (7.5mg fenbendazole/kg bodyweight). Over 650kg dose rate of 15mL per 50kg bodyweight. Dose the mob to the heaviest animal by liveweight. Do not underdose. A representative sample of animals should be weighed before treatment.

PRECAUTIONS

When used as directed no specific precautions are required other than those normally associated with the careful handling of stock especially weak or pregnant animals. Internal parasites may develop resistance to drenches (anthelmintics). If drench (anthelminitic) failure with this preparation is suspected, seek veterinary confirmation and notify the manufacturer.

Liveweight (kg)	Dose Volume (mL)	Doses per 1L Pack
50	15	66
100	30	33
150	45	22
200	60	16
250	75	13
300	90	11
350	105	9
400	120	8
450	135	7
500	150	6
550	165	6
600	180	5
650	195	5

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT (SHEEP, GOATS, CATTLE): DO NOT USE less than 14 days before slaughter for human consumption

MILK (CATTLE): NIL.

MILK (GOATS): Milk collected from goats within 24 hours ([2] milkings) following treatment MUST NOT BE USED for human consumption or processing, or fed to kids.

MILK (SHEEP): DO NOT USE on lactating or pregnant ewes where milk or milk products may be used for human consumption.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): Sheep and Cattle: DO NOT USE less than 14 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Intervet/Schering-Plough Animal Health on 1 800 226 511 or the APVMA website (www.apvma.gov.au/residues/ESI.shtml).

EXPORT SLAUGHTER INTERVAL (ESI): Goats: An ESI has not been

established. For advice on the ESI contact Intervet/Schering-Plough Animal Health on 1 800 226 511 before using this product.

GENERAL INSTRUCTIONS

COOPERS PANACUR 25 does not stain wool or skin or predispose animals to photosensitisation. COOPERS PANACUR 25 is palatable and readily accepted by livestock. COOPERS PANACUR 25 is active against both mature (adult) and immature (larval) nematode parasites. COOPERS PANACUR 25 has activity (80% to 90%) against inhibited (hypobiotic) larvae of *Ostertagia* in cattle. COOPERS PANACUR 25 is ready to use and non-clogging. COOPERS PANACUR 25 may be administered concurrently with vaccines, jetting and dipping operations and pour-on lice treatments. COOPERS PANACUR 25 has ovicidal activity in sheep and goats, reducing worm egg hatching. For this to be of benefit, sheep and goats should not be placed on clean pasture until 12 hours have elapsed after drenching. Resistance may develop to any drench and it is advisable that a drench resistance test be conducted before any drench is used. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance.

COOPERS®
PANACUR® 25
ORAL ANTHELMINTIC FOR SHEEP, CATTLE AND GOATS

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Ph 13 11 26.

Additional Information is listed in the Material Safety Data Sheet.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

Dispose of empty container by wrapping with paper and putting in garbage.

Warranty

Intervet Australia Pty Ltd (IAPL) warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by any act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

SPECIALIST ADVICE IN EMERGENCY ONLY

Intervet/Schering-Plough Animal Health

1800 226 511

ALL HOURS – AUSTRALIA WIDE

COOPERS ANIMAL HEALTH

A division of Intervet Australia Pty Ltd
91–105 Harpin Street, Bendigo East, Vic 3550

STORE below 30°C (Room Temperature). Do not freeze. Store in tightly closed original container in a well-ventilated area. Do not store for prolonged periods in direct sunlight.

APVMA Approval No. 37097/1L/0109

FOR BATCH AND EXPIRY SEE BELOW

- © Registered Trademarks
- © Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.
- © 2009

1L