



VyLyte

CONCENTRATED ORAL REHYDRATION LIQUID FOR LIVESTOCK & PETS

VyLyte is a non-antibiotic oral solution for use as an aid to replace lost fluids and electrolytes during periods of stress. Suitable for piglets, pigs, calves, cattle, lambs, sheep, goats, horses, foals, dogs & cats.

ACTIVE CONSTITUENTS PER LITRE

GLUCOSE	278.75g
SODIUM CHLORIDE	53.44g
GLYCINE	38.56g
POTASSIUM PHOSPHATE MONOBASIC	25.53g
CITRIC ACID MONOHYDRATE	3.28g
POTASSIUM CITRATE	0.75g

- ✓ Prevents and treats mild cases of scouring
- ✓ Suitable for a variety of species
- ✓ Aids in the prevention of dehydration by replacing electrolytes and lost liquids
- ✓ Formulated with a balanced mix of glucose, glycine, citrate, and electrolytes
- ✓ Can be used to treat pregnancy toxemia in ewes
- ✓ Helps to improve dogs and cats' health suffering from viral diseases
- ✓ Fights against acidosis



Contact Vetsense:



RATE OF USE

Vetsense VyLyte is a liquid concentrate requiring dilution with fresh clean water where final volume is 12.5 times quantity of concentrate. To make up 1L of ready to consume liquid, measure 80mL of VyLyte and add water to make 1L. (measure 10mL and add water to 125mL, measure 160mL and add water to make 2L, etc.)

DILUTION GUIDE

VYLYTE LIQUID	WATER	TOTAL VOLUME
10mL	115mL	125mL
20mL	230mL	250mL
40mL	460mL	500mL
80mL	920mL	1000mL
160mL	1840mL	2000mL

Each VyLyte pack size makes:
250mL = 3.125L, 1L = 12.5L, 5L = 62.5L, 20L = 250L

ALWAYS ENSURE CLEAN DRINKING WATER IS AVAILABLE WHEN USING VETSENSE VYLYTE. DISCARD SOLUTION DAILY AND REPLACE WITH FRESH SOLUTION EVERY 24 HOURS.

Provide diluted VETSENSE VyLyte at first sign of nutritional scouring to aid and replace lost fluid and electrolytes.

SEEK VETERINARY ADVICE IF TREATING SICK AND DEBILITATED ANIMALS

PIGS

Administer in a clean drinking vessel as soon as signs of scouring appear.

Suckling Piglets: A litter of week-old piglets will consume roughly 2-3L/day. Check that diluted solution, drinking water or sow's milk is always available. Check quantity of diluted solution every 12 hours and replenish as necessary. Piglets can be dosed directly using an appropriate applicator; 20-30mL diluted VyLyte solution, 3 or 4 times a day.

Weaned Pigs: Limit feed intake for 1-2 days, continue supply of drinking water with weaners consuming 1-2L/day/animal as a guide.

Treatment period (all pigs): Diluted VyLyte solution can be given for 4 days. If scouring persists, application is suitable for 8 days in total.

CALVES

Bought-in Calves: At the first feed on arrival, feed 2L of diluted VyLyte solution instead of milk. At the next feed, give 1L of diluted VyLyte solution but do not mix with milk or milk replacer. Feed normal diet post application.

Scouring Calves: 1L to 1.5L of diluted VyLyte solution per 10kg of bodyweight daily into two or more daily doses. Severe dehydration can happen if scouring is well established, increase diluted solution over 3 or 4 separate doses. Directions: To retain nutrient and energy consumption, continue normal milk feeding during periods of scouring. Fresh water should always be available, with diluted VyLyte solution given between milk feeds. Replace solution every 24 hours.

Scouring Lambs/Kids

Give 150mL-200mL of fresh diluted VyLyte solution, 2-4 times daily warming solution to body temperature and given through a feeding bottle with teat.

Pregnancy Toxaemia in EWES/GOATS

Give 160mL undiluted VyLyte solution per animal using applicable drench gun. Repeat every 4-6 hours if required or as directed by a veterinary surgeon.

FOALS

For scouring or dehydration, give 1-2L of fresh diluted VyLyte 2-3 times a day or as directed by a veterinary surgeon.

✓ **Available Pack Sizes:** 250mL, 1L, 5L, 20L

✓ **Withholding Periods:** Nil (0 days)

✓ **Store below 30°C (room temperature)**

DOGS & CATS DOSAGE GUIDE

WEIGHT	DILUTED VYLYTE SOLUTION	FREQUENCY
Cats & Dogs under 5kg	125mL	2-3 times/day
Dogs 5-10kg	250mL	2-3 times/day
Dogs 11-20kg	500mL	2-3 times/day
Dogs 20kg+	750mL	2-3 times/day

